

12

EUROPEAN PATENT APPLICATION

21 Application number: 89401845.6

51 Int. Cl.⁵: **C 07 H 19/167**
A 61 K 31/70

22 Date of filing: 28.06.89

30 Priority: 21.07.88 EP 88401896

43 Date of publication of application:
14.03.90 Bulletin 90/11

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **MERRELL DOW PHARMACEUTICALS INC.**
2110 East Galbraith Road
Cincinnati Ohio 45215-6300 (US)

72 Inventor: **Casara, Patrick**
12, rue de la Scierie
F-67117 Ittenheim (FR)

Danzin, Charles
18, rue Geller
F-67000 Strasbourg (FR)

74 Representative: **Gillard, Marie-Louise et al**
Cabinet Beau de Loménie 55, Rue d'Amsterdam
F-75008 Paris (FR)


54 **Novel S-adenosyl methionine decarboxylase inhibitors.**

57 This invention relates to novel chemical compounds useful as S-adenosylmethionine decarboxylase inhibitors, to the processes useful for their preparation and to their use in the treatment of a variety of condition and disease states associated with a rapid proliferation of cell growth.


EP 0 358 536 A2

This invention relates to novel chemical compounds useful as S-adenosylmethionine decarboxylase inhibitors, to the processes useful for their preparation and to their use in the treatment of a variety of condition and disease states associated with a rapid proliferation of cell growth.




More specifically, this invention relates to compounds of the formula




and the pharmaceutically acceptable salts thereof, wherein Ad represents adenosinyl, R is hydrogen or a C₁₋₇ alkyl, and Q represents moieties of formulae Ia to Ie, said formulae being depicted as:



la

**1b****lc**

Id



le


with m being zero or one, q being zero or one with the proviso that the sum of m and q is less than 2,

V is H or -COOH,

W is H, F, Cl or Br,


Z is H, F, Cl or Br,

each X and each Y being H or F,




-C≡CH or -CH_{3-n}F_n with n being 1, 2 or 3.

Of course, the "H₂N-" moiety (which is attached to Q in formula I) is attached to the carbon atom bearing the "V" substituent in each of formulae 1a to 1e (e.g.,



wher in Q is represented by lb).

The "Ad" moiety (i.e., the adenosinyl moiety comprised of 1H-purine-6-amine attached to β -D-ribofuranosyl) has the structure




In those instances wherein Q is representative of formula 1a, it is to be noted that only one of the moieties may be one for any given compound. In those instances wherein R is other than H, the C₁₋₇ alkyl moiety is preferably methyl and ethyl but all the straight, branched-chain and cyclized manifestations are included with methyl being preferred and ethyl, propyl, isopropyl, butyl, t-butyl, pentyl, hexyl, cyclohexyl, methyl and the like being especially included. In those instances wherein Q is representative of formula 1e, it is preferred that said compounds be in their *cis*-configuration rather than their *trans*-configuration. In those instances wherein Q is representative of formula 1d the mono-, di-, tri- and tetrafluoro substituted moieties (as well as the unsubstituted moieties) are contemplated. Preferred compounds are those of formula I wherein Q is 1e, are those wherein W and Z are H and V is H or COOH, and A is methyl or ethyl.

Illustrative examples of pharmaceutically acceptable salts of the compounds of this invention are those formed with inorganic acids preferably with hydrochloric, hydrobromic, sulfuric or phosphoric acids and with organic acids such as methane sulfonate, salicylic, maleic, malonic, tartaric, citric and ascorbic acids. These salts may be prepared by standard techniques and procedures well known in the art.


In essence, the preparation of the compounds of formula may be effected by techniques and chemical processes analogously known in the art; the choice of the specific route being dependent upon the usual factors in pharmaceutical research institutions such as availability and cost of starting materials, time and difficulties in separation and purification of intermediates and final compounds and such other factors well known and generally appreciated by those of ordinary skill in the art.

In general the preparation of the compounds of formula may be depicted by the reaction sequence of reaction sequence of reaction scheme A, as follows:


Reaction Scheme A




wherein Ad' is adeninyl of the formula




10 R_2X' is a reactant of the following formulae (3a to 3f).




3a




3b




3c



3d



3e



3f

50 wherein Pg is an N-protecting group, preferably t-butoxycarbonyl (Boc) or phthalimido (Pht) (in which case, of course, the H of the PgNH moiety is not present) and m, n, q, R, R_1 , V, W, X, Y and Z are as defined in Formula 1, and X' is OTf (triflate) or chloro, bromo or iodo, (R_2 , of course, being the moieties of 3a to 3f attached to the X' moiety), with the exception that in appropriate cases in may be a reaction-protection derivative of the COOH function, preferably a t-butoxy derivative, and Ac is an acyl moiety, preferably acetate, and R_3 is t-butyl.


55 In effecting the condensation of reactants 2 and 3 when X' represents a halide conditions A-a are utilized wherein equimolar quantities of the reactants are reacted together in the presence of a base, (preferably potassium carbonate), in a basic solvent, (preferably acetonitrile), at temperatures of about 30°C to 80°C. When conditions A-b are utilized, i.e., when X' is a triflate, the reactants are heated together at about 30°C to 80°C in the presence of a base, (preferably triethylamine), in a basic solvent, (preferably dimethylformamide).
60 Removal of the N-protecting groups is readily effected by standard techniques, e.g., treatment with 1N sulfuric acid at room temperature for 24-48 hours followed by treatment with an alcohol (preferably ethanol) at about 0°C when the protecting group is t-butoxycarbonyl, and when the protecting group is phthalimido, removal is effected using an ethanolic solution of a hydrazine (using classical techniques) the latter being used when R_2 contains a fluoro atom. Removal of the isopropylidene protecting group of the ribofuranosyl moiety is easily
65 effected by hydrolysis at room temperatures, (preferably using 1N sulfuric acid), generally simultaneously with

the N-protecting groups. Isolation and purification of the intermediate and final products of reaction scheme A is effected by standard techniques, e.g., recrystallization, HPLC, flash chromatography (on silica gel) and the like.

The preparation of the intermediates required for the condensation of reaction scheme A, i.e., those intermediates defined for R_2X' , may be effected by the use of analogously known procedures such as those outlined in the below described generic processes which are illustrated in the below particularized examples.

In those instances wherein R_2X' represents subgeneric group 3e, the reaction proceeds under A-a conditions wherein X' is preferably chloro and the N-protecting group is Boc, the appropriate W, W, Z-substituted-N-protected-4-chloro-2-butene-1-amine may conveniently be prepared by the following reaction scheme.

Reaction Scheme B



wherein the protecting groups (Pht and Boc) and V, W and Z are as previously defined and (THP) is tetrahydropyran.

In step B-a the diol is reacted with dihydropyran in the presence of catalytic quantities of pyridinium-p-toluene sulfonate at about 0°C in an anhydrous solvent (or mixture) (e.g., $\text{CH}_2\text{Cl}_2:\text{THF}$; 2:1) for about 24-48 hours. Conversion of B-2 to B-3 is initiated by a Mitsunobu-type intermolecular dehydration reaction on treatment with diethylazodicarboxylate (DEAD) and triphenylphosphine under mild neutral conditions under an inert atmosphere (nitrogen) at about 0°C in an anhydrous solvent (e.g., THF) in the presence of phthalimide; the reaction continuing at room temperature for about 12 hours.


The resulting B-3 products, upon treatment with hydrazine hydrate in ethanol at reflux for about 12 hours, to remove the phthalimido and THP protecting groups and the free amine is re-protected with di-tert-butylidicarbonate by refluxing in dichloromethane. The alcohols (B-4) are converted to their chlorides by reaction with mesylchloride under basic conditions (TEA) in an anhydrous solvent, preferably dichloromethane. These cis products of Formula 3e, after purification, generally using flash chromatographic techniques on silica gel, are ready for condensation with the reactants of Formula 2, according to the techniques described for reaction scheme A.

In those instances wherein it is desired to prepare the trans configuration of compounds of 3-e, it is preferred to utilize a W, Z-V-substituted N-protected trans-1-bromo-4-amino-2-butene, (i.e., 3-e), the reactants are readily prepared by reacting a V-W, Z-substituted trans-1-bromo-4-amino-2-buten with


potassium phthalimide in anhydrous DMF at about 50°C for 24 hours according to standard procedures well known in the art. The necessary R₂X' reactants of the class 3-c are readily prepared from the appropriate W, Z, V₂-substituted α,α -dichloroxylene wherein the compound is subjected to a displacement reaction with potassium phthalimide to form an α -phthalimido- α' -chloroxylene by heating the reactants at about 50°C for about 24 hours in anhydrous DMF and the so-formed compound is purified by the usual techniques of flash chromatography from silica gel. Starting from the appropriately V, W, Z-substituted 3-chloro-2-chloromethyl-1-propene the desired R₂X' reactants of class 3-a may similarly be prepared by the foregoing described displacement reaction with potassium phthalimide by heating the reactants at about 50°C for about 24 hours in anhydrous dimethylfluoromethane followed by purification with the usual techniques, e.g., flash chromatography. In those instances wherein the particular V, W, Z-substituted reactant is not a known compound, such compounds may be prepared by techniques and procedures well understood and known in the art.

In addition to the specific examples described below, chemistry for the preparation of cis-5'-(4-amino-4-carboxy-2-butenyl)methyladenosine may be analogously derived from Tolman and Sedmera's article (Tetrahedron Letters, Vol. 29, No. 47, pp. 6183-6184, 1988) "Unsaturated Amino Acids: Synthesis of Trans-3,4-Didehydro Analogues of L-Ornithine and L-Arginine". The application of this chemistry is schematically represented by the following reaction scheme.


Reaction Scheme C




(6)




(7)




(8)




(9)



(10)



(11)



(12)

In effecting the foregoing reaction scheme step (a) involves the dibromination of (6) by reaction with bromine, said reaction being placed in a suitable solvent (e.g., CCl_4) at room temperature. The resulting dibromo analog is debrominated by reaction with potassium t-butoxide in tetrahydrofuran or with an amine such as DBU. The so-obtained compound (7) is sequentially treated with (1) trifluoroacetic acid at 25°C for 20 minutes, (2) treated with thionyl chloride at 25°C for 3 hours, and (3) treated with DiBAL in tetrahydrofuran at -30°C for 1 hour to produce compound (8). Step (c) involves the sequential treatment of (8) with a base (e.g., $\text{NaOH}/\text{H}_2\text{O}$ in tetrahydrofuran for 20 minutes, followed by treatment with diluted HCl at 50°C to produce

compound (9). This compound is treated with isobutylene, in the presence of catalytic amounts of sulfuric acid and the resulting alcohol is converted to its corresponding chloride by treatment with mesyl chloride to produce compound (10). This compound is then subjected to reaction with the adenosine derivatives of formula (2) according to the procedure of Reaction Scheme A (wherein compound (10) corresponds to R₂X' with X' being chloro) to produce a compound analogously corresponding to compounds 4 [i.e., compound (11)].

The resulting triple-bond-containing compound is partially reduced using hydrogenation in the presence of a Lindlar catalyst (H₂/PdSO₄) and the resulting butene is treated with sulfuric acid (to remove the t-butoxide and isopropylidene protecting groups). The final step is to subject the so-produced penultimate compound to acylase I (Merck) at a pH of 7.2 at 37°C to remove the N-protecting acyl moiety to produce a desired compound (12), e.g., cis-5'-deoxy-5'-(4-amino-4-carboxy-2-butene)methylaminoadenosine.

The following examples illustrate the preparation of the necessary intermediates and final products of this invention.

EXAMPLE 1

Preparation of CIS-5'-DEOXY-5'-(4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

Step A:

CIS-4-TETRAHYDROPYRANYLOXY-2-BUTENE-1-OL

Dihydropyran (9.1 ml, 100 mmol) was added dropwise to a cooled (0°C) solution of 2-butene-1,4-diol (8.8 g, 10 mmol) and pyridinium paratoluenesulfonate (0.25 g, 10 mmol) in anhydrous dichloromethane:tetrahydrofuran (2:1). The mixture was stirred two days at 0°C then concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (ethyl acetate:hexane 3:7) to give 8.3 g of the title compound (49%).

Step B:

CIS-1-PHTHALIMIDO-4-TETRAHYDROPYRANYLOXY-2-BUTENE

Under a nitrogen atmosphere diethylazodicarboxylate (1.6 ml, 10 mmol) was added to a cooled (0°C) solution of cis-4-tetrahydropyranyloxy-2-butene-1-ol (1.7 g, 10 mmol), triphenyl phosphine (2.2 g, 10 mmol) and phthalimide (1.47 g, 10 mmol) in anhydrous tetrahydrofuran (50 ml). When the addition was completed (5 min) the reaction mixture was allowed to warm at room temperature and was stirred 12 h. Then the mixture was concentrated *in vacuo*, diluted with ethyl acetate (200 ml) and washed with brine (150 ml). After usual work-up (the aqueous phase was extracted three times with 100 ml portions of ethyl acetate), the organic phase was dried over magnesium sulfate, filtered and concentrated *in vacuo* the product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 2:8) to give 1.9 g of the title compound (64%).

Step C:

CIS-TERTIOBUTOXYCARBONYL-4-HYDROXY-2-BUTENYL-1-AMINE

A solution of cis-1-phthalimido-4-tetrahydropyranyloxy-2-butene (1.9 g, 6.3 mmol) and hydrazine hydrate (0.35 ml, 6.9 mmol) in ethanol (20 ml) was heated under reflux 12 hours. Then the mixture was concentrated *in vacuo*, diluted with 1N hydrochloric acid (20 ml) and heated under reflux for two hours. Then the phthalylhydrazide was filtered off and the filtrate was concentrated *in vacuo*. The residue was taken in dichloromethane (100 ml) neutralized with triethylamine (pH 8.9) and a solution of di-tert-butylidicarbonate (1.65 g, 7.5 mmol) in dichloromethane (5 ml) was added. The mixture was heated under reflux overnight and, after usual work-up, the product was obtained by flash chromatography on silica gel (ethyl acetate: hexane; 25:75) (0.8 g, 74%).

Step D:

CIS-N-TERTIOBUTOXYCARBONYL-4-CHLORO-2-BUTENYL-1-AMINE

Mesyl chloride (0.6 ml, 7.6 mmol) was added to a cooled (0°C) solution of cis-tert-butoxycarbonyl-4-hydroxy-2-butenyl-1-amine (1.3 g, 7 mmol) and triethylamine (1.1 ml, 7.6 mmol) in anhydrous dichloromethane (30 ml). The mixture was stirred overnight and, after usual work-up, the title product was purified by flash chromatography on silica gel (ethyl acetate: hexane; 2:8) (0.8 g, 57%).

Step E:

CIS-5'-DEOXY-5'

(N-TERTIOBUTOXYCARBONYL-4-AMINO-2-BUTENYL)-METHYL-AMINO-2',3'-ISOPROPYLIDENEADENOSINE

A solution of cis-N-tertiobutoxycarbonyl-4-chloro-2-butenyl-1-amin (0.6 g, 3 mmol), 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (0.97 g, 3 mmol), potassium carbonate (0.42 g, 3 mmol) and sodium iodide (0.05 g, 0.3 mmol) in acetonitrile (20 ml) was heated under reflux overnight. Then the mixture was diluted with ethyl acetate, washed with brine and dried over magnesium sulphate. Then the product was purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) (1.1 g, 55%).

Step F:

CIS-5'-DEOXY-5'-(4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

A solution of cis-5'-deoxy-5'[(N-tertiobutoxycarbonyl-4-amino-2-butenyl)methyl-amino]-2',3'-isopropylideneadenosine (0.9 g, 1.8 mmol) in 1N sulphuric acid (5 ml) was left two days at room temperature. Then the mixture was diluted with ethanol (200 ml) and cooled (0°C) overnight. The precipitate was filtered off, dissolved in the minimum amount of water and then re-precipitated with ethanol (200 ml). This procedure was repeated twice to give the title compound (0.5 g), mp: 260°C decomposed.

EXAMPLE II

Preparation of TRANS-5'-DEOXY-5'-(4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

Step A:

TRANS-1-BROMO-4-PHTHALIMIDO-2-BUTENE

A mixture of trans-1,4-dibromo-2-butene (6.4 g, 30 mmol) and potassium phthalimide (5.6 g, 30 mmol) in anhydrous dimethyl formamide (200 ml) was heated at 50°C for 24 h. Then the reaction mixture was concentrated in vacuo, dissolved in ethyl acetate, washed with brine and the pure title product was obtained by flash chromatography on silica gel (ethyl acetate: hexane; 15:85) (3.2 g, 40%).

Step B:

TRANS-5'-DEOXY-5'-(4-PHTHALIMIDO-2-BUTENYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of trans-1-bromo-4-phthalimido-2-butene (2 g, 7.5 mmol), potassium carbonate (1.6 g, 11.5 mmol) and 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (2.4 g, 7.5 mmol) in anhydrous acetonitrile (100 ml) was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, dissolved in dichloromethane, filtered and purified by flash chromatography on silica gel (chloroform: diethylamine; 98:2) to afford the title compound (1.25 g, 33%).

Step C:

TRANS-5'-DEOXY-5'-(4-TERTIOBUTOXYCARBONYLAMINO-3-BUTENYL)-METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of trans-5'-deoxy-5'-(4-phthalimido-2-butenyl)methylamino-2',3'-isopropylideneadenosine (1 g, 2 mmol) and hydrazine hydrate (0.1 ml, 2 mmol) in absolute ethanol was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, dissolved in water (30 ml) and the pH was adjusted to 4 with glacial acetic acid and cooled to 0°C. Then the mixture was filtered off and the filtrate neutralized with triethylamine to pH 9 and concentrated *in vacuo*. Then the residue was dissolved in dichloromethane, and ditertibutylidicarbonate (0.45 g, 2 mmol) was added. The mixture was heated under reflux overnight and, after usual work-up, the product was purified by flash chromatography on silica gel (diethylamine:dichloromethane; 2:98) to give the title compound (0.5 g, 51%).

Step D:

TRANS-5'-DEOXY-5'-(4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

A suspension of trans-5'-deoxy-5'-(4-tertiobutoxycarbonyl-amino-2-butenyl)methylamino-2',3'-isopropylideneadenosine (0.4 g, 0.96 mmol) in 1N sulphuric acid (3 ml) was stirred 2 days at room temperature. Then the mixture was diluted with absolute ethanol (100 ml) and cooled at 0°C overnight. The product was filtered off, dissolved in the minimum amount of water and precipitated with ethanol (100 ml). This procedure was repeated twice to afford the title compound (0.16 g). mp: 250-260°C decomposed.

EXAMPLE III

Preparation of 5'-DEOXY-5'-(4-AMINO-2-BUTYNYL)METHYLAMINOADENOSINEStep A:5 1-CHLORO-4-PHTHALIMIDO-2-BUTYNE

A mixture of 1,3-dichloro-2-butyne (4.9 ml, 50 mmol) and potassium phthalimide (5.6 g, 30 mmol) was heated at 50°C during 24 h. Then the mixture was concentrated *in vacuo*, diluted with ethyl acetate and, after usual work-up, the product was purified by flash chromatography on silica gel to give 4.3 g of the title compound (62%).

10

Step B:5'-DEOXY-5'-(4-PHTHALIMIDO-2-BUTYNYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

15 A mixture of 1-chloro-4-phthalimido-2-butyne (1.4 g, 6 mmol), 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.6 g, 5 mmol) and sodium iodide (0.075 g, 0.5 mmol) in anhydrous acetonitrile (100 ml) was heated under reflux overnight. Then the mixture was concentrated, diluted with dichloromethane, filtered and purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) to give the title compound (1.6 g, 64%).

20 Step C:5'-DEOXY-5'-(4-TERTIOBUTOXYCARBONYLAMINO-2-BUTYNYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

25 A mixture of 5'-deoxy-5'-(4-phthalimido-2-butyne)methylamino-2',3'-isopropylideneadenosine (1 g, 1.9 mmol) and methyl hydrazine (0.5 ml, 10 mmol) in absolute ethanol (3 ml) was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, dissolved in a mixture of tetrahydrofuran:water (1:1, 200 ml), and a solution of di-tert-butyl dicarbonate (0.5 g, 2.5 mmol) in tetrahydrofuran (10 ml) was added. The pH of the mixture was adjusted to 9 with triethylamine and then the mixture was heated under reflux for 24 h. Then the reaction mixture was concentrated *in vacuo*, diluted with ethyl acetate and, after usual work-up, the product was obtained by flash chromatography on silica gel (diethylamine:chloroform; 2:98) (0.5 g, 56%).

30

Step D:5'DEOXY-5'-(4-AMINO-2-BUTYNYL)METHYLAMINOADENOSINE

35 A suspension of 5'-deoxy-5'-(4-tert-butoxycarbonylamino-2-butyne)methylamino-2',3'-isopropylideneadenosine (0.4 g, 0.82 mmol) in 1N sulphuric acid (25 ml) was stirred 2 days at room temperature. Then the mixture was diluted with ethanol (100 ml) and stirred at 0°C overnight. The product was filtered off, dissolved in the minimum amount of water and diluted with ethanol (100 ml). This procedure was repeated twice to afford pure 5'-deoxy-5'-(4-amino-2-butyne)methylaminoadenosine as white crystals (0.2 g). mp: 230-240°C decomposed. This compound, of course, can be reduced to form the corresponding cis double-bonded compound.

40

EXAMPLE IV

45

Preparation of 5'-DEOXY-5'-(ORTHO-AMINOMETEYL BENZYL)METHYLAMINOADENOSINEStep A:

50

 α -PHTHALIMIDO- α' -CHLOROXYLENE

A mixture of α,α' -dichloroxylylene (8.75 g, 50 mmol) and potassium phthalimide (5.6 g, 30 mmol) was heated to 50°C for 24 h. Then the reaction mixture was concentrated *in vacuo*, dissolved in ethyl acetate and, after usual work-up, the desired compound was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 15:85) (6 g, 65%).

55

Step B:5'-DEOXY-5'-(ORTHO-PHTHALIMIDO-METHYLBENZYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

60

A mixture of α -phthalimido- α' -chloroxylylene (1.6 g, 5.5 mmol), potassium carbonate (0.7 g, 5 mmol) sodium iodide (0.07 g, 0.5 mmol) and 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.5 g, 4.7 mmol) in anhydrous acetonitrile was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, dissolved in dichloromethane, filtered and then purified by flash chromatography on silica gel (chloroform:diethylamine 98:2) to give the title compound (1.8 g, 67%).

65

Step C:5'-DEOXY-5'-(ORTHO-TERTIOBUTOXYCARBONYLAMINOMETHYLBENZYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of 5'-deoxy-5'-(ortho-phthalimido-methylbenzyl)methylamino-2',3'-isopropylideneadenosine (1.3 g, 2.3 mmol) and hydrazine hydrate (0.12 ml, 2.3 mmol) in absolute ethanol (100 ml) was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, diluted in water (30 ml), and glacial acetic acid was added to adjust at pH 4 and left at 0°C. Then the mixture was filtered off and the filtrate was neutralized with triethylamine to adjust the pH of the reaction mixture around 9. Then the mixture was concentrated *in vacuo*, diluted with dichloromethane, and ditertiobutyldicarbonate (0.5 g, 2.3 mmol) was added. Then the mixture was heated under reflux overnight and, after usual work-up, the title compound (0.8 g, 67%) was isolated by flash chromatography on silica gel (chloroform: diethylamine; 98:2).

Step D:5'-DEOXY-5'-ORTHO-AMINOMETHYLBENZYL)METHYLAMINOADENOSINE

A suspension of 5'-deoxy-5'-(ortho-tertobutoxycarbonyl aminomethylbenzyl)methylamino-2',3'-isopropylideneadenosine (0.45 g, 0.83 mmol) in 1N sulphuric acid (25 ml) was stirred two days at room temperature. Then the mixture was diluted with ethanol (100 ml) and stored at 0°C overnight. The precipitate was filtered off, dissolved in the minimum amount of water and reprecipitated with ethanol (100 ml). This procedure was repeated twice to give the title compound (0.4 g). mp: 230-240°C decomposed.

EXAMPLE V5'-DEOXY-5'-(3-AMINO-2-METHYLENEPROPYL)METHYLAMINOADENOSINEStep A:1-PHTHALIMIDO-3-CHLORO-2-METHYLENEPROPANE

A mixture of 3-chloro-2-chloromethyl-1-propene (6.55 g, 50 mmol) and potassium phthalimide (5.6 g, 30 mmol) in anhydrous dimethylformamide (200 ml) was heated two days at 50°C. Then the mixture was concentrated *in vacuo* and, after usual work-up, the product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 15:85) (4.2 g, 78%).

Step B:5'-DEOXY-5'-(3-PHTHALIMIDO-2-METHYLENEPROPYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of 1-phthalimido-3-chloro-2-methylenepropene (0.87 g, 5 mmol), potassium carbonate (0.7 g, 5 mmol), sodium iodide (0.08 g, 0.5 mmol) and 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.6 g, 5 mmol) in anhydrous acetonitrile (100 ml) was heated two days under reflux. Then the mixture was concentrated *in vacuo*, diluted with dichloromethane, filtered and the product was purified by flash chromatography on silica gel (diethyl amine: chloroform; 2:98) to give 2.85 g (78%) of the title compound.

Step C:5'-DEOXY-5'-(3-TERTIOBUTOXYCARBONYLAMINO-2-METHYLENEPROPYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of 5'-deoxy-5'-(3-phthalimido-2-methylenepropyl)methylamino-2',3'-isopropylideneadenosine (2.3 g, 4.4 mmol), methyl hydrazine (1.5 ml, 30 mmol) in absolute ethanol (5 ml) was heated two days under reflux. Then the mixture was concentrated *in vacuo*, dissolved in chloroform (5 ml), the pH was adjusted around 9 with triethylamine and then a solution of ditertiobutyl dicarbonate (8.8g, 4.4 mmol) in chloroform (5 ml) was added. The resulting mixture was heated overnight under reflux and, after usual work-up, the product was purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) to give 1.25 g (64%) of the title compound.

Step D:5'-DEOXY-5'-(3-AMINO-2-METHYLENEPROPYL)METHYLAMINOADENOSINE

A suspension of 5'-deoxy-5'-(3-tertobutoxycarbonylamino-2-methylenepropyl)methylamino-2',3'-isopropylideneadenosine (0.65 g, 1.3 mmol) in 1N sulphuric acid (4 ml) was stirred two days at room temperature. Then the mixture was diluted with absolute ethanol (150 ml) and left at 0°C overnight. The precipitate was filtered off,

dissolved in a minimum amount of water and diluted with absolute ethanol (150 ml). This procedure was repeated twice to afford the title compound as white crystals (0.55 g, mp: 230-240°C decomposed).

5

EXAMPLE VIPreparation of 5'-DEOXY-5'-(4-AMINO-2,2-DIFLUOROBUTYL)METHYLAMINOADENOSINE10 Step A:4-PHTHALIMIDO-2,2-DIFLUOROBUTYL-TRIFLUOROMETHANESULFONATE

Triflic anhydride (1.1 ml, 6.6 mmol) was added to a cooled (0°C) solution of 4-phthalimido-2,2-difluoro-1-butanol (1.53 g, 6 mmol), pyridine (0.53 ml, 6.6 mmol) in anhydrous dichloromethane (50 ml). The mixture was stirred 1 h at 0°C and, after usual work-up, the product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 20:80) to give 1.8 g (78%) of the title compound.

Step B:20 5'-DEOXY-5'-(4-PHTHALIMIDO-2,2-DIFLUOROBUTYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of 4-phthalimido-2,2-difluorobutyl-trifluoromethanesulfonate (1.8 g, 4.6 mmol), 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.3 g, 4.3 mmol) and triethylamine (0.6 ml, 4.3 mmol) in anhydrous dimethylformamide was heated two days at 50°C. Then the mixture was concentrated *in vacuo* and the product was purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) (1.7 g, 70%).

Step C:25 5'-DEOXY-5'-(4-TERTIOBUTOXYCARBONYLAMINO-2,2-DIFLUOROBUTYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A mixture of 5'-deoxy-5'-(4-phthalimido-2,2-difluorobutyl)methylamino-2',3'-isopropylideneadenosine (1.5 g, 2.7 mmol) and hydrazine hydrate (0.135 g, 2.7 mmol) in ethanol (20 ml) was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, diluted with water, and glacial acetic acid was added until the pH was adjusted to 4. The mixture was left at 0°C and then filtered off. The filtrate was neutralized to pH 9 with triethylamine, concentrated *in vacuo*, diluted with dichloromethane and then di-tert-butylidicarbonate (0.6 g, 2.7 mmol) was added. The mixture was heated under reflux overnight and, after usual work-up, the product was purified by flash chromatography on silica gel (diethylamine: chloroform; 2:98) to give 1.1 g (75%) of the title compound.

40 Step D:5'-DEOXY-5'-(4-AMINO-2,2-DIFLUOROBUTYL)METHYLAMINOADENOSINE

A suspension of 5'-deoxy-5'-(4-tert-butoxycarbonylamino-2,2-difluorobutyl)methylamino-2',3'-isopropylideneadenosine in 1N sulphuric acid (4.5 ml) was stirred two days at room temperature. Then the mixture was diluted with ethanol (100 ml) and left overnight at 0°C. The precipitate was filtered off, dissolved in a minimum amount of water and precipitated with ethanol (150 ml). This procedure was repeated twice to afford the title compound (0.5 g, 60%) as white crystals (mp: 240°C decomposed).

50

EXAMPLE VIIPreparation of CIS-5'-DEOXY-5'-(4-AMINO-2-FLUORO-2-BUTENYL)METHYLAMINO ■ ADENOSINE55 Step A:CIS-4-PHTHALIMIDO-2-FLUORO-1-TETRAHYDOPYRANYL-2-BUTENE

A mixture of cis-4-chloro-2-fluoro-1-tetrahydropyranyl-2-buten (6.3 g, 30 mmol) and potassium phthalimide (5.6 g, 30 mmol) in anhydrous dimethyl formamide (200 ml) was heated at 50°C for 24 h. Then the reaction mixture was concentrated *in vacuo*, dissolved in ethyl acetate, washed with brine and the pure title compound cis-4-phthalimido-2-fluoro-2-tetrahydropyranyl-2-butene (6 g, 70%) was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 2:8).

65

Step B:CIS-N-TERTIOBUTOXYCARBONYL-2-FLUORO-4-HYDROXY-2-BUTENYL-1-AMINE

A solution of cis-4-phthalimido-2-fluoro-2-tetrahydro pyran-2-yl-2-butene (5.7 g, 20 mmol) and hydrazine hydrate (1.1 ml, 22 mmol) in ethanol (30 ml) was heated under reflux for 12 h. Then the mixture was concentrated *in vacuo*, diluted with 1N HCl (20 ml) and heated under reflux for 2 h. Then the phthalhydrazide was filtered off and the filtrate was concentrated *in vacuo*. The residue was taken up in dichloromethane (150 ml), neutralized with triethylamine until pH 9, and a solution of ditertibutyldicarbonate (5 g, 22 mmol) in dichloromethane (10 ml) was added. The mixture was heated under reflux overnight and, after usual work-up, the product was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 25:75) (3 g, 75%).

Step C:CIS-N-TERTIOBUTOXYCARBONYL-2-FLUORO-4-CHLORO-2-BUTENYL-1-AMINE

Mesylic chloride (0.9 ml, 11 mmol) was added to a cold (0°C) solution of cis-N-tertiobutoxycarbonyl-2-fluoro-4-hydroxy-3-butenyl-1-amine (2.05 g, 10 mmol) and triethylamine (1.6 ml, 11 mmol) in anhydrous dichloromethane (40 ml). The mixture was stirred overnight and, after usual work-up, the title compound cis-N-tertiobutoxycarbonyl-2-fluoro-4-chloro-2-butenyl-1-amine was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 15:85) (1.7 g, 75%).

Step D:CIS-5-DEOXY-5'-(4-TERTIOBUTOXYCARBONYLAMINO-2-FLUORO-2-BUTENYL)METHYLAMINO-2',3'-ISO-PROPYLIDENEADENOSINE

A solution of 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.65 g, 5 mmol), cis-N-tertiobutoxycarbonyl-2-fluoro-4-chloro-2-butenyl-1-amine (1.2 g, 4 mmol), potassium carbonate (0.7 g, 4 mmol) and sodium iodide (0.07 g, 0.5 mmol) in anhydrous acetonitrile (30 ml) was heated under reflux overnight. The mixture was concentrated *in vacuo*, diluted with ethyl acetate, washed with brine and dried over MgSO₄. The product was purified by flash chromatography on silica gel (diethylamine: chloroform; 2:98) (1.7 g, 70%).

Step E:CIS-5'-DEOXY-5'-(4-AMINO-2-FLUORO-2-BUTENYL)METHYLAMINOADENOSINE

A suspension of cis-5'-deoxy-5'-(4-tertiobutoxycarbonylamino-2-fluoro-2-butenyl)methylamino-2',3'-isopropylideneadenosine in 1N sulphuric acid (5 ml) was stirred for 2 days at room temperature. Then the mixture was diluted with absolute ethanol (200 ml) and kept at 0°C overnight. The precipitate was collected, dissolved in a minimum of water, and reprecipitated with absolute ethanol (200 ml). This procedure was repeated twice to give the title compound cis-5'-deoxy-5'-(4-amino-2-fluoro-2-butenyl)methylaminoadenosine (1 g, 75%; mp: 250-260°C decomposed).

EXAMPLE VIIIPreparation of 5'-DEOXY-5'-(3'-AMINO-2,2-DIFLUOROPROPYL)METHYLAMINOADENOSINEStep A:ETHYL 2,2-DIFLUORO-3-HYDROXYPROPIONATE

A mixture of paraformaldehyde (4.5 g, 50 mmol), ethyl difluorobromoacetate (10.2 g, 50 mmol) and activated zinc dust (3.3 g, 40 mmol) in anhydrous tetrahydrofuran was heated under reflux for 0.5 h. Then the mixture was treated with a saturated aqueous solution of ammonium chloride and extracted with diethyl ether. After usual work-up the desired compound ethyl 2,2-difluoro-3-hydroxypropionate was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 25:75) (4.1 g, 53%).

Step B:ETHYL 2,2-DIFLUORO-3-TETRAHYDROPYRANYLOXYPROPIONATE

Dihydropyran (2 ml, 22 mmol) was added to a solution of ethyl 2,2-difluoro-3-hydroxypropionate (3.1 g, 20 mmol) and pyridinium p-toluenesulfonate (0.25 g, 1 mmol) in anhydrous dichloromethane (50 ml). The mixture was stirred overnight at room temperature and the desired compound ethyl 2,2-difluoro-3-tetrahydropyranyloxypropionate was obtained by flash chromatography on silica gel (ethyl acetate:hexane; 15:85) (4 g, 80%).

Step C:2,2-DIFLUORO-3-TETRAHYDROPYRANYLOXY-1-PROPANOL

5 A solution of ethyl 2,2-difluoro-3-tetrahydropyranyloxypropionate (3.5 g, 15 mmol) in absolute ethanol (10 ml) was added dropwise to a slurry of sodium borohydride (0.57 g, 15 mmol) at room temperature in absolute ethanol (20 ml). Then the mixture was stirred an additional hour at room temperature. Then the mixture was concentrated *in vacuo*, hydrolyzed with aqueous ammonium chloride, extracted with ethyl acetate and dried over magnesium sulfate. The product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 25:75) (2.7 g, 90%).

Step D:2,2-DIFLUORO-3-TETRAHYDROPYRANYLOXYPROPYL TRIFLUOROMETHANESULFONATE

15 Triflic anhydride (1.8 ml, 11 mmol) was added to a cold (0°C) solution of 2,2-difluoro-3-tetrahydropyranyloxy-1-propanol (9.6 g, 10 mmol), pyridine (0.9 ml, 11 mmol) in anhydrous dichloromethane (50 ml). The mixture was stirred 1 h at 0°C and, after usual work-up, the product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 15:85) (2.6 g, 80%).

Step E:2,2-DIFLUORO-3-PHTHALIMIDO-1-TETRAHYDROPYRANYLOXYPROPANE

20 A mixture of 2,2-difluoro-3-tetrahydropyranyloxypropyl trifluoromethanesulfonate (2.3 g, 7 mmol), potassium phthalimide (1.4 g, 7.7 mmol) and anhydrous dimethylformamide (50 ml) under nitrogen was stirred and heated at 85°C overnight. After cooling, salts are filtered off, and the solvent was removed *in vacuo*. The residue was taken up in dichloromethane (100 ml), washed with 0.5M NaOH (30 ml) and brine. The organic phase was separated, dried over magnesium sulfate and concentrated. The desired compound 2,2-difluoro-3-phthalimido-1-tetrahydropyranyloxypropane was purified by flash chromatography on silica gel (ethyl acetate:hexane; 20:80) (2 g, 90%).

Step F:2,2-DIFLUORO-3-PHTHALIMIDO-1-PROPANOL

30 A solution of 2,2-difluoro-3-phthalimido-1-tetrahydropyranyloxypropane (2 g, 6.15 mmol), paratoluene sulfonic acid (0.1 g) in absolute ethanol was stirred overnight at room temperature. Then the mixture was concentrated *in vacuo*, diluted with ethyl acetate and washed with brine. The organic phase was separated, dried over magnesium sulfate and concentrated *in vacuo*. The crude alcohol 2,2-difluoro-3-phthalimido-1-propanol (1.4 g) was used for the next step without further purification.

Step G:2,2-DIFLUORO-3-PHTHALIMIDO-PROPYL TRIFLUOROMETHANE SULFONATE

40 Triflic anhydride (1.1 ml, 6.6 mmol) was added to a cold (0°C) solution of 2,2-difluoro-3-phthalimido-1-propanol (1.4 g, 6 mmol), pyridine (0.5 ml, 6.6 mmol) in anhydrous dichloromethane (30 ml). The mixture was stirred 1 h at 0°C and, after usual work-up, the product was purified by flash chromatography on silica gel (ethyl acetate:hexane; 20:80) (1.7 g, 75%).

Step H:5'-DEOXY-5'-(2,2-DIFLUORO-3-PHTHALIMIDO-PROPYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

50 A mixture of 2,2-difluoro-3-phthalimido-propyl trifluoromethane sulfonate (1.5 g, 4 mmol), 5'-deoxy-5'-methylamino-2',3'-isopropylideneadenosine (1.2 g, 4.2 mmol) and triethylamine (0.55 ml, 4.2 mmol) in anhydrous dimethyl formamide was heated 2 days at 50°C. Then the mixture was concentrated *in vacuo* and the product was purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) (1.5 g, 75%).

Step I:5'-DEOXY-5'-(2,2-DIFLUORO-3-TERTIOBUTOXYCARBONYLAMINOPROPYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

60 A mixture of 5'-deoxy-5'-(2,2-difluoro-3-phthalimidopropyl)methylamino-2',3'-isopropylideneadenosine (1.1 g, 2 mmol) in ethanol (10 ml) was heated under reflux overnight. Then the mixture was concentrated *in vacuo*, diluted with 1N acetic acid until pH 4 was reached, and cooled at 0°C. The precipitate was filtered off and the filtrate was neutralized until pH 9 with triethylamine and concentrated *in vacuo*. The residue was taken up in dichloromethane and di-tert-butyl dicarbonate (0.45 g, 2 mmol) was added. The mixture was heated under

reflux overnight and, after usual work-up, the product was purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98) (0.8 g, 70%).

Step J:

5'-DEOXY-5'-(3-AMINO-2,2-DIFLUOROPROPYL)METHYLAMINOADENOSINE

A suspension of 5'-deoxy-5'-(2,2-difluoro-3-tertiobutoxycarbonylamino-2',3'-isopropylideneadenosine (0.8 g, 1.5 mmol) in 1N sulphuric acid (4 ml) was stirred 2 days at room temperature. Then the mixture was diluted with absolute ethanol (150 ml) and kept at 0°C overnight. The precipitate was collected, dissolved in a minimum of water, and reprecipitated with absolute ethanol (150 ml). This procedure was repeated twice to give the title compound 5'-deoxy-5'-(3-amino-2,2-difluoropropyl)methylaminoadenosine (0.6 g, 80%: mp: 250-260°C decomposed).

EXAMPLE IX

Preparation of CIS-5'-DEOXY-5'-(4-CARBOXY-4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

Step A:

2-AMINO-5-HYDROXY-3-PENTYNOIC ACID

A mixture of glyoxylic acid monohydrate (23 g, 250 mmol), propargyl alcohol (16.8 g, 300 mmol), copper (II) chloride (3.2 g, 25 mmol) and ammonium acetate (49 g, 600 mmol) in ethanol (100 ml) is heated under reflux for 6 h. Then the reaction mixture is concentrated *in vacuo*, diluted with water (50 ml), acidified to pH 5 with 1N HCl and washed twice with ether (100 ml). Then the aqueous solution is poured on an ion exchange resin column (DOWEX 50, H⁺). The column is eluted with 1 M ammonium hydroxide to give the title compound 2-amino-5-hydroxy-3-pentynoic acid.

Step B:

TERTIOBUTYL-2-AMINO-5-HYDROXY-3-PENTYNOATE

A suspension of 2-amino-5-hydroxy-3-pentynoic acid (12.5 g, 100 mmol) concentrated in sulphuric acid (2 ml) and isopropylene (50 ml) in a sealed Parr's flask is shaken 2 days at room temperature. The crude product, after evaporation of the excess of isopropylene, is used for the next step without further purification.

Step C:

TERTIOBUTYL-2-TERTIOBUTOXYCARBONYLAMINO-5-HYDROXY-3-PENTYNOATE

A solution of the crude tertibutyl-2-amino-5-hydroxy-3-pentynoate (100 mmol), ditertibutyldicarbonate (22 g, 100 mmol) and triethylamine (25 ml, 200 mmol) in chloroform is heated under reflux overnight. Then, after usual work-up, the product is purified by flash chromatography on silica gel (ethyl acetate:hexane; 20:80).

Step D:

CIS-TERTIOBUTYL-2-TERTIOBUTOXYCARBONYLAMINO-5-HYDROXY-3-PENTENOATE

A solution of tertibutyl-2-tertiobutoxycarbonylamino-5-hydroxy-3-pentynoate (13.6 g, 50 mmol) in ethanol (200 ml) is hydrogenated in presence of Lindlar catalyst (0.6 g) at atmospheric pressure and room temperature. In 3 h one equivalent of hydrogen (1.1 liters) is taken up. Then the catalyst is removed by filtration and the mixture is concentrated *in vacuo* which will yield a clear oil. The title compound is obtained by flash chromatography on silica gel (ethyl acetate:hexane; 15:85).

Step E:

CIS-TERTIOBUTYL-2-TERTIOBUTOXYCARBONYLAMINO-5-CHLORO-3-PENTENOATE

Mesyl chloride (0.9 ml, 11 mmol) is added to a cold (0°C) solution of cis-tertibutyl-2-tertiobutoxycarbonylamino-5-hydroxy-3-pentenoate (2.75 g, 10 mmol) and triethylamine (1.6 ml, 11 mmol) in anhydrous dichloromethane (50 ml). The mixture is stirred overnight and, after usual work-up, the title compound is purified by flash chromatography on silica gel (ethyl acetate:hexane; 20:80).

Step F:

CIS-5'-DEOXY-5'-(4-TERTIOBUTOXYCARBONYL-3-TERTIOBUTOXYCARBONYLAMINO-2-BUTENYL)METHYLAMINO-2',3'-ISOPROPYLIDENEADENOSINE

A solution of cis-tertibutyl-2-tertiobutoxycarbonylamino-5-chloro-3-pentenoate (1.5 g, 5 mmol), 5'-deoxy-

5'-methylamino-2',3'-isopropylideneadenosine (1.6 g, 5 mmol), potassium carbonate (0.7 g, 5 mmol) and sodium iodide (0.8 g, 0.5 mol) in acetonitrile (30 ml) is heated under reflux overnight. After usual work-up, the product is purified by flash chromatography on silica gel (diethylamine:chloroform; 2:98).

5 Step G:

CIS-5'-DEOXY-5'-(4-CARBOXY-4-AMINO-2-BUTENYL)METHYLAMINOADENOSINE

A suspension of cis-5'-deoxy-5'-(4-tert-butoxycarbonyl-3-tert-butoxycarbonylamino-2-butenyl)methylamino-2',3'-isopropylideneadenosine (1.5 g, 3 mmol) in 1N sulphuric acid (5 ml) is stirred 2 days at room temperature. Then the mixture is diluted with ethanol (200 ml) and kept at 0°C overnight. The precipitate is collected, dissolved in a minimum amount of water, and reprecipitated with ethanol (200 ml). This procedure is repeated twice and will yield the title compound cis-5'-deoxy-5'-(4-carboxy-4-amino-2-butenyl)methylaminoadenosine.

(Usual work-up involves the extraction of the product from the aqueous phase by three extractions with the organic solvent (as in Step C, Example I) and the organic phase dried over magnesium sulfate, filtered off and concentrated *in vacuo*.)

The compounds of Formula I are inhibitors of decarboxylase enzymes which are involved in polyamine formation and therefore such compounds are useful as pharmacological agents. In particular, the compounds of Formula I are potent and irreversible inhibitors of S-adenosylmethionine decarboxylase (Ado Met DC) and therefore significantly interfere with the formation of spermine and spermidine and thus are useful adducts in the armamentarium of researchers and clinicians in the study of polyamine formation and in the treatment of conditions and diseases related to the rapid proliferation of cell growth. Of particular importance is the use of the compounds of this invention in conjunction with known ornithine decarboxylase inhibitors, known antitumoral agents and with immunomodulators known for their use in diseases associated with the rapid proliferation of normal and transformed cells.

As is well known in the art, polyamines are associated with both normal and rapid proliferation of cells and, as is also well known, the levels of polyamines are high in embryonic systems, the testes and patients suffering from diseases associated with rapid proliferation of cell growth. It is also known that there is a correlation between the activity of the decarboxylase enzymes of ornithine, S-adenosylmethionine, arginine and lysine and polyamine formation. Thus with this interrelationship the compounds of this invention, by their unique ability to inhibit S-adenosylmethionine decarboxylase, are also useful to study the biochemical and pharmacological consequences of polyamine biosynthesis blockade in mammals, plants, bacteria and protozoa.

More specifically, some of the more promising end-use applications of the compounds of this invention are the following applications.

The use of an Ado Met DC inhibitor of this invention, alone but more effectively in conjunction with an ornithine decarboxylase (ODC) inhibitor in mammals as postcoital contraceptives, inducers of menstruation, and as first-trimester abortifacients is clear for such use does not surgically invade the cavity of the uterus, does not require hospitalization, and can be administered with the minimum of medical supervision.

As is well known the use of single agents in anti-neoplastic chemotherapy has been only marginally successful and limited to a very few malignancies. Thus treatment of such malignancies as leukemia (e.g., acute lymphocytic leukemia, Hodgkins' disease), melanoma and metastatic melanoma, multiple myeloma, bladder carcinoma, prostate carcinoma (as well as benign renal adenocarcinoma, gastric adenocarcinoma, prostatic hypertrophy), pancreatic adenocarcinoma, colon cancer, lung carcinoma, fibrosarcoma and mammary carcinoma will be enhanced by the use of the Ado Met DC inhibitors of this invention in conjunctive therapeutic regimens with prior art drug combinations. For example, the use of the instant Ado Met DC inhibitors, preferably with ODC inhibitors and/or with interferon will significantly facilitate a lowering of untoward side effects and increase survival time when used in conjunction with such well-known drug combinations as vincristine, amethopterin, mercaptopurine, prednisone (VAMP), cyclophosphamide, arabinosylcytosine, oncovin (COAP), mechlorethamine, procarbazine (MOPP), ABVD, CHOP, CAF, FAM and PVE (see review by Berger, N.A. (1986), J. Clin. Invest. (1131-1135). Additionally the use of the compounds of this invention improves the efficacy of the prevention of metastasis in cancers of the breast, colon and bladder with ODC inhibitors. Another specific clinical aspect of the compounds of this invention is the enhancement of the synergism between ODC inhibitors and α -interferon in the treatment of cancers, particularly metastatic malignant melanoma. Still another aspect of the enhancement of the unique interaction between ODC inhibitors and methylglyoxal bisguanyldihydrazone (MGBG; methyl GAG), particularly in cases with recurrent primary brain tumors (e.g., astrocytomas). Still further, the compounds of this invention may be useful in the treatment of hyperproliferative disorders such as psoriasis.

In addition to the foregoing, the compounds of this invention may be used to treat diseases which are caused by infections with animal parasites, particularly parasitic protozoa and parasitic nematodes. In the treatment of diseases caused by these parasites, the compounds of this invention may be used alone, or in combination with ornithine decarboxylase inhibitors and/or in combination with other agents known to be useful in the treatment of such diseases. In some instances, such as in treating Chagas Disease, it is preferred to utilize arginine decarboxylase inhibitors in conjunction with the compounds of this invention. Of particular interest is the treatment of African trypanosomiasis, Chagas Disease and *Pneumocystis carinii* pneumonia (PCP) in patients suffering from AIDS (particularly in conjunction with an ODC inhibitor), cryptosporidiosis and

malaria.

Still more specifically, the compounds of this invention are particularly useful in treating:

(1) diseases caused by *Onchocerca Volvulus*, a filarial nematode living in subcutaneous tissue causing skin lesions and eye lesions (river blindness). These diseases are commonly treated with diethylcarbamazine which kills the microfilariae, but not the adult worms;

(2) diseases caused by *Wuchereria bancrofti*, a thread-like nematode the adults of which live in thin lymphatic vessels and which cause lymphangitis, dermatitis and cellulitis. These diseases are commonly treated with diethylcarbamazine but this treatment is known to be inadequate;

(3) diseases caused by *Loa loa*, a filarial worm causing Loaiasis characterized by hot erythematous Calabar swelling on extremities and periorbital tissues which also have been treated with diethylcarbamazine;

(4) diseases caused by *Trichomonas vaginalis* a sexually transmitted flagellate protozoa causing trichomoniasis which is commonly treated with metronidazole;

(5) diseases caused by *Giardia lamblia*, a flagellated protozoan parasite causing giardiasis in domestic dogs and wild animals as well as man, for which the drugs of choice are quinacrine hydrochloride and metronidazole;

(6) diseases caused by *Toxoplasma gondii*, a protozoan parasite of the sub-class coccidia causing congenital toxoplasmosis which may be treated with pyrimethamine and sulfadiazine;

(7) diseases caused by the malaras of the genus *Plasmodium*, e. g., *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium ovale* and *Plasmodium malariae* which are the causes of malaria; the usual treatment is with chloroquine, quinine sulfate, pyrimethamine and sulfadiazine. In the treatment of this disease it is better to utilize the compounds of this invention in conjunctive therapy with the foregoing or with ODC inhibitors;

(8) diseases caused by *Trypanosoma cruzi*, a protozoa causing Chagas Disease, a disease which has a history of being difficult to treat. In the treatment of this disease with the compounds of this invention it is recommended that they be used in conjunctive therapy with arginine and agmatine decarboxylase inhibitors;

(9) diseases caused by the African trypanosomes, *Trypanosoma brucei gambiense* and *Trypanosoma brucei rhodesiense*, two subspecies of hemoflagellates which are responsible for African Trypanosomiasis, a disease treated by suramin and more recently by eflornithine. Of particular use in the treatment of this disease are the specific compounds cis-5'-deoxy-5'-(4-amino-2-butenyl)methylaminoadenosine and cis-5'-deoxy-5'-(4-carboxy-4-amino-2-butenyl)methylaminoadenosine.

(10) diseases caused by *Leishmania tropica* and *Leishmania mexicana* which are responsible for cutaneous leishmaniasis, and visceral leishmaniasis (also called kala azar or black fever) which is caused by *Leishmania donovani*.

Furthermore, an Ado Met DC inhibitor of this invention may be used (alone or in combination with ODC inhibitors) as anti-infective agents being effective in the control of bacteria, fungi and viruses which are dependent upon polyamines for growth, for example *E. coli*, *Enterobacter*, *H. influenzae*, *Mycobacteria* species, *Staphylococcus aureus*, *Klebsiella*, viruses such as poxviruses, Herpes viruses, picornaviruses and influenza viruses. In the use of the compounds of this invention it is preferred to use such ODC inhibitors α -difluoromethylornithine, α -monofluoromethylornithine, α -ethylornithine, (c)-2-(fluoromethyl)dehydroornithine (and the methyl, ethyl and other esters thereof), and (2R,5R)-6-heptyne-2,5-diamine; said compounds and their uses being adequately described as to their preparation, their ODC inhibitory properties and to their end-use applications (see Inhibition of Polyamine Metabolism, (1987) edited by McCann, P.P., Pegg, A.E., and Sjoerdsma, A.).




The S-adenosylmethionine decarboxylase inhibitory properties of the compounds of Formula I may readily be determined by standard laboratory procedures well known in the art. For example, cis-5'-deoxy-5'-(4-amino-2-butenyl)methylaminoadenosine produces inactivation of rat liver Ado Met Dc *in vitro* with a $t_{1/2}$ = 16 minutes at 0.1 μ M, a $t_{1/2}$ = 1.6 minutes at 1 μ M and a $t_{1/2}$ = 0.8 minutes at 2 μ M.

As pharmacologically useful agents the compounds of Formula I can be administered in various manners to the patient being treated to achieve the desired effect. The compounds can be administered alone or in combination with another according to standard techniques for conjunctive therapy, bearing in mind that the compounds of this invention preferably enhance established protocols when used to treat neoplasms. The compounds are preferably administered in the form of a pharmaceutical preparation. In general, the compounds may be administered orally, parenterally, for example, intravenously, intraperitoneally, or subcutaneously, infusionally or topically, as determined by factors well-known and appreciated by the skilled artisan. The amount of compound administered will vary over a wide range and can be any effective amount, depending on the patient to be treated, the condition being treated and the mode of administration. The effective amount of compound administered will vary from about 0.2 mg/kg to 200 mg/kg of body weight of the patient per treatment dose and preferably will be about 1 mg/kg to about 50 mg/kg of body weight of the patient per treatment dose.

The solid unit dosage forms can be of the conventional type. Thus, the solid form can be a capsule which can be of the ordinary gelatin type containing a novel compound of this invention and a carrier, for example, lubricant and inert fillers, such as lactose, sucrose and corn starch. In another embodiment, the novel compounds are tableted with conventional tablet bases such as lactose, sucrose or corn starch in

10 The compounds can be administered in the form of a depot injection or implant preparation which may be formulated in such a manner as to permit a sustained release of the active ingredient. The active ingredients can be compressed into pellets or small cylinders and implanted subcutaneously or intramuscularly as depot injections or implants. Implants may employ inert materials such as biodegradable polymers or synthetic silicones, for example, Silastic, silicone rubber manufactured by the Dow-Corning Corporation.

25 Claims

$$\begin{array}{c} \text{W} \quad \text{Z} \\ \diagdown \quad \diagup \\ \text{C} \\ \parallel \\ \text{H}_2\text{N}-\text{CH}(\text{V})-(\text{CH}_2)_m-\text{C}-(\text{CH}_2)_q\text{CH}_2- \end{array} \quad \text{Ia}$$



BNSDOCID: <EP__0358536A2_1_>

W is H, F, Cl or Br,
 Z is H, F, Cl or Br,
 each X and each Y being H or F,






R_1 is $-\text{CH}=\text{C}\begin{smallmatrix} \text{W} \\ \text{Z} \end{smallmatrix}$, $-\text{CH}=\text{C}=\text{C}\begin{smallmatrix} \text{W} \\ \text{Z} \end{smallmatrix}$,

$-\text{C}\equiv\text{CH}$ or $-\text{CH}_3\text{F}_n$ with n being 1, 2 or 3.


2. A compound of Claim 1 wherein R is methyl or H.
3. A compound of Claim 1 wherein Q represents a moiety of formula 1e.
4. A compound of Claim 2 wherein the compound is in its cis-configuration.
5. A compound of Claim 1 wherein Q represents a moiety of formula 1a.
6. A compound of Claim 1 wherein Q represents a moiety of formula 1c.
7. A compound of Claim 3, said compound being cis-5'-deoxy-5'-(4-amino-2-butenyl)methylaminoadenosine.
8. A compound of Claim 3, said compound being cis-5'-deoxy-5'-(4-amino-2-butenyl)aminoadenosine.
9. A compound of Claim 3, said compound being cis-5'-deoxy-5'-(4-amino-4-carboxy-2-butenyl)methylaminoadenosine.
10. A compound of Claim 3, said compound being cis-5'-deoxy-5'-(4-amino-4-carboxy-2-butenyl)aminoadenosine.
11. A compound of Claim 1, said compound being 5'-deoxy-5'-(3-amino-2-methylenepropyl)methylaminoadenosine.
12. A compound of Claim 1, said compound being 5'-deoxy-5'-(3-amino-2-methylenepropyl)aminoadenosine.
13. A compound of the formula



wherein R_2 is a moiety of the following formulae


13a13b13c13d13e

with m being zero or one, q being zero or one, with the proviso that the sum of m and q is less than 2,
 V' is H, COOH or a reaction-protected COOH,
 W is H, F, Cl or Br,
 Z is H, F, Cl or Br,
 each X and each Y being H or F,




$-\text{C}\equiv\text{CH}$ or $-\text{CH}_3\text{-F}_n$ with n being 1, 2 or 3.


14. A process for preparing a compound of the formula




and the pharmaceutically acceptable salts thereof, wherein Ad represents adenosinyl, R is hydrogen or a C₁₋₇ alkyl, and Q represents moieties of formulae 1a to 1e, said formulae being:




1a




1b



1c




1d




1e







with m being zero or one, q being zero or one with the proviso that the sum of m and q is less than 2,
 V is H or -COOH,
 W is H, F, Cl or Br,
 Z is H, F, Cl or Br,
 each X and each Y being H or F,



-C \equiv CH or -CH_{3-n}F_n with n being 1, 2 or 3;
 which comprises reacting a compound of the formula



with a R₂X' reactants, R₂X' being a compound of the formulae

3a3b3c3d3e3f

wherein Pg is an N-protecting moiety, V' is H, COOH or COOR₃ with R₃ being a reaction protection moiety, and X' is a triflate, chloro, bromo or iodo, and Ac is an acyl moiety, preferably acetate, said reaction being effected with equimolar quantities of the involved reactants in the presence of a base, preferably in a basic solvent, at temperatures of about 30°C to 80°C, followed by the removal of any protecting groups using standard techniques, except that when a reactant of formula 3f is utilized the removal of the nitrogen, carboxyl and/or hydroxy protecting groups attached to the condensation reaction products is effected after the chemical reduction of the butyne moiety to its corresponding butene moiety by hydrogenation in the presence of a Lindlar catalyst, and optionally converting the so-produced compounds to their pharmaceutically acceptable salts.



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 358 536 A3**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89401845.6

(51) Int. Cl.⁵: C07H 19/167, A61K 31/70

(22) Date of filing: 28.06.89

(30) Priority: 21.07.88 EP 88401896

(43) Date of publication of application:
14.03.90 Bulletin 90/11

(54) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(88) Date of deferred publication of the search report:
03.04.91 Bulletin 91/14

(71) Applicant: MERRELL DOW
PHARMACEUTICALS INC.
2110 East Galbraith Road
Cincinnati Ohio 45215-6300(US)

(72) Inventor: Casara, Patrick
12, rue de la Scierie
F-67117 Ittenheim(FR)
Inventor: Danzin, Charles
18, rue Geiler
F-67000 Strasbourg(FR)

(74) Representative: Gillard, Marie-Louise et al
Cabinet Beau de Loménie 55, Rue
d'Amsterdam
F-75008 Paris(FR)

(54) Novel S-adenosyl methionine decarboxylase inhibitors.

(57) This invention relates to novel chemical compounds useful as S-adenosylmethionine decarboxylase inhibitors, to the processes useful for their preparation and to their use in the treatment of a variety of condition and disease states associated with a rapid proliferation of cell growth.

EP 0 358 536 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 40 1845

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. CL5)
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 19, no. 5, May 1976, pages 684-691; C.-D. CHANG et al.: "Analogues of S-adenosylhomocysteine as potential inhibitors of biological transmethylation. Synthesis of analogues with modifications at the 5'-thioether linkage" * Abstract; page 684, compound 1d; page 685, scheme II; page 687, table 1 and figure 1 * -- -- --	1,2,6,13, 14	C 07 H 19/167 A 61 K 31/70
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 23, no. 2, February 1980, pages 121-127, American Chemical Society; M. PANKASKIE et al.: "Inhibitors of polyamine biosynthesis. 8. Irreversible inhibition of mammalian S-adenosyl-L-methionine decarboxylase by substrate analogues" * Abstract; page 122, Scheme II, compound 21; page 123, figure 1 * -- -- --	1,2	
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 25, no. 5, May 1982, pages 550-556, American Chemical Society; M. KOLB et al.: "Synthesis and biochemical properties of chemically stable product analogues of the reaction catalyzed by S-adenosyl-L-methionine decarboxylase" * Abstract; page 551, Scheme II, compounds 7,8; page 552, compounds 10,13; page 553, figure 1 and table 1 * -- -- --	1,2,6,13, 14	
X	CHEMICAL ABSTRACTS, vol. 100, no. 13, 26th March 1984, pages 698-699, abstract no. 103834n, Columbus, Ohio, US; M. DAVIS et al.: "The synthesis of the N-methyl analog of S-adenosylmethionine: NMR observation of diastereomers", & AUST. J. CHEM. 1983, 36(8), 1623-7 * Abstract * -- -- -- -- / --	1,2,13,14	
The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		18 January 91	SCOTT J.R.M.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ----- & : member of the same patent family, corresponding document			



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 89 40 1845

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	BIOCHEMISTRY, vol. 21, no. 7, 30th March 1982, pages 1535-1541, American Chemical Society; C.S.G. PUGH et al.: "Effects of S-adenosylhomocysteine analogues on vaccinia viral messenger ribonucleic acid synthesis and methylation" * Abstract; page 1536, table 1, especially AdoDab; page 1538, table II *	1,2	
X,P	JOURNAL OF ORGANIC CHEMISTRY, vol. 53, no. 21, 14th October 1988, pages 4952-4961, American Chemical Society; A.A. MINNICK et al.: "General synthetic approach to stable nitrogen analogues of S-adenosylmethionine" * The whole document *	1,2,13,14	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.5)
Place of search		Date of completion of search	Examiner
The Hague		18 January 91	SCOTT J.R.M.
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone</p> <p>Y: particularly relevant if combined with another document of the same category</p> <p>A: technological background</p> <p>O: non-written disclosure</p> <p>P: intermediate document</p> <p>T: theory or principle underlying the invention</p> <p>E: earlier patent document, but published on, or after the filing date</p> <p>D: document cited in the application</p> <p>L: document cited for other reasons</p> <p>8: member of the same patent family, corresponding document</p>			

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.